arXiv:nucl-ex/0205017v1 25 May 2002

# Flow with PMD: Past and Future[1]

Sudhir Raniwala

Physics Department, University of Rajasthan, Jaipur 302004, India

**Abstract**

Measurements of azimuthal distribution of inclusive photons using the fine granularity preshower photon multiplicity detector (PMD) at CERN SPS are used to obtain anisotropy in the azimuthal distributions. These results are used to estimate the anisotropy in the neutral pion distributions. The results are compared with results of charged particle data, both for first order and second order anisotropy. Assuming the same anisotropy for charged and neutral pions, the anisotropy in photons is estimated and compared with the measured anisotropy. The effect of neutral pion decay on the correlation between the first order and the second order event plane is also discussed. Data from PMD can also be used to estimate the reaction plane for studying any anisotropy in particle emission characteristics in the ALICE experiment at the Large Hadron Collider. In particular, we show that using the event plane from the PMD, it will be possible to measure the anisotropy in J/$\psi$ absorption (if any) in the ALICE experiment.

[1]Invited talk at the Fourth International Conference on Physics and Astrophysics of Quark Gluon Plasma, 26-30 Nov.2001, Jaipur, India

# 1 Introduction

A preshower photon multiplicity detector (PMD) was first used in the WA93 experiment to measure the inclusive photon yield and their spatial distributions to study S+Au collisions at SPS energies [1]. The measurement of the azimuthal distributions of these photons resulted in reporting the first ever observation of anisotropy at these energies [2]. Part of the observed anisotropy was due to correlations from the neutral pion decay. The observed anisotropy in photons could not be used to determine the anisotropy in neutral pions, nor could the quantitative contribution from decay correlations be determined at that time. However, the observed anisotropy was found to be greater than what was observed in a similar sample of VENUS events, which implicitly include the effect of energy momentum correlations in neutral pion decay.

There now exist methods that enable a determination of the neutral pion anisotropy from a measurement of the anisotropy in azimuthal distribution of inclusive photons [3]. In the present paper we shall show the results on anisotropy in photon emission and its centrality dependence. We shall use the method detailed in [3] to estimate the anisotropy in neutral pions and compare the results with the anisotropy measured in the charged pions in the same pseudorapidity region. The effect of finite granularity of the detector(s) and the effect of efficiency and purity of the data sample will also be investigated.

The measures of anisotropy are the coefficients in the Fourier expansion of the azimuthal angle distribution. The precision of these coefficients systematically increases for increasing anisotropy and for increasing multiplicity. The measured azimuthal angle distribution of the photons in the ALICE environment can also provide an estimate of the reaction plane of the event. The possibility of measuring any dependence of particle emission characteristics on the event plane direction is investigated, in particular the possible anisotropic absorption of J/$\psi$ resonance [4].

# 2 WA98 Experiment and DATA

The preshower PMD measured the inclusive distribution of photons in Pb+Pb collisions on an event by event basis in the WA98 experiment in the pseudorapidity region 2.8 to 4.2 [5]. The Silicon Pad Multiplicity Detector (SPMD) measured charged particles in the region 2.35 to 3.75. The present analysis deals with the data in the region 3.25 to 3.75 corresponding to 100% azimuthal coverage for the two detectors in the region of overlap. The basic features of the PMD, the minimum bias multiplicity distribution and the pseudorapidity distributions can be seen in [6]. Details of data reduction, efficiency and purity

of the detected samples can be seen in [5]. The analysis has been carried out for the seven centrality classes. The total number of events in every centrality class and the corresponding percentage of cross section are shown in Table 1.

Table 1: WA98 DATA

| $E_T$ Cut | Percentage of Cross Section | Number of Events |
|---|---|---|
| $> 347.6$ | 0 - 5 % | 68 K |
| 298.6 - 347.6 | 5 - 10 % | 75 K |
| 225.5 - 298.6 | 10 - 20 % | 54 K |
| 170.2 - 225.5 | 20 - 30 % | 18 K |
| 124.4 - 170.2 | 30 - 40 % | 19 K |
| 89.9 - 124.3 | 40 - 50 % | 18 K |
| 40.0 - 89.9 | 50 - 80 % | 37 K |

# 3 Analysis and Results

For every event, the event planes of both orders are determined using the relation

$$\psi_n = \frac{1}{n}\left(\tan^{-1}\frac{\Sigma w_i \sin n\phi_i}{\Sigma w_i \cos n\phi_i}\right) \tag{1}$$

The $\phi_i$ are the azimuthal angles of the emitted particles and the $w_i$ are the weight factors. For the azimuthal distribution of the number of particles, these weight factors are all equal to one. The data have been corrected for acceptance. The azimuthal distribution of the inclusive photons (and of charged particles) is plotted for every centrality bin and the inverse of occupancies have been taken as weights in obtaining the event plane angles in the equation above. This has been done for both PMD and SPMD for all centralities. A typical distribution for the second order event plane angles is shown in Fig. 1a for the $\gamma$-like particles recorded in the PMD, both before and after the acceptance correction. $\gamma$-like particles include the inclusive photons and some charged particles which deposit an energy comparable to energy deposited by a photon [5].The flatness of the $\psi_2$ distribution shows that the detector acceptance effects have been corrected for.

The distribution of the azimuthal angles of the emitted particles with respect to the event planes enables a determination of the coefficients $v_1$ and $v_2$




(a) (b)

Figure 1: (a) $\psi_2$ distribution of $\gamma$-like hits for various centralities before and after the acceptance correction factors (b) Azimuthal angle distribution of particles with respect to the second order (corrected) event plane angles. First column is for $\gamma$-like in the PMD and the second column is for the charged particles in the SPMD

characterising directed and elliptic flow. Both these coefficients are determined in more than one way, differing in details and is a good recipe for consistency checks.

We determine

1. $v_1' = \langle \cos(\phi - \psi_1^{est}) \rangle$ : a measure of the directed flow in the direction of $\psi_1^{est}$, where the average is over all particles of all events.
2. $v_2' = \langle \cos 2(\phi - \psi_2^{est}) \rangle$ : a measure of the ellipticity about $\psi_2^{est}$ where the average is again over all particles of all the events.

The distribution of the azimuthal angles of the particles with respect to the event plane is shown in Fig. 1b for the second order, both for PMD and for SPMD, for all centralities. To avoid auto correlations, the event plane has been obtained from all the particles excluding the particle in question. The $v_n'$ values are then determined by fitting this distribution to

$$r(\phi) = \frac{1}{2\pi} \left[ 1 + 2v_1' \cos(\phi - \psi_1) + 2v_2' \cos 2(\phi - \psi_2) \right] \tag{2}$$

Distribution of particle azimuthal angles is also obtained with respect to the acceptance uncorrected event plane angle. Similar distribution is obtained

for mixed events. The effect of detector anisotropy cancels in the ratio of the two distributions and is used to determine the $v_n$' values. The $v'_n$ values obtained this way are the same as obtained for the case by fitting the distribution to Fig. 1b. The prime on the $v_n$ indicates that these are projected on the estimated reaction plane and not the true reaction plane and therefore need a correction factor.

## 3.1 Event Plane Resolution Correction

Since the estimated event plane fluctuates about the actual event plane due to finite particle multiplicity, the average correction factor for the event plane determination, termed the resolution correction factor (RCF) is determined. Experimentally, RCF is obtained using the subevent method described in [7]. Here every event is divided randomly into two subevents of equal multiplicity and the angle $\psi_n$ is determined for each subevent. This enables a determination of a parameter $\chi_n$ directly from the experimental data using the relation [7, 8]:

$$\frac{N_{events}(n|\psi_n^a - \psi_n^b| > \pi/2)}{N_{total}} = \frac{e^{-\frac{\chi_n^2}{4}}}{2} \tag{3}$$

where $N_{total}$ denotes the total number of events, $\psi_n^a$, $\psi_n^b$ are the estimated angles of the two subevents (labelled $a$ and $b$) and the numerator on the left denotes the number of events having the angle between subevents greater than $\pi/2n$. The parameter $\chi_n$ is then used to determine RCF according to the relation given in the Ref. [7].

For the data sample used in the present analysis, the resolution correction factors are shown in Fig. 2. $RCF_n$ is the correction factor for anisotropy of the order n and $RCF_{12}$ is the correction factor required when one determines the second order anisotropy with respect to the event plane of the first order. Smaller values of these factors introduce larger errors on the estimated values of the anisotropy.

## 3.2 Anisotropy Values

The resultant values of $v_n$ for the PMD and for the SPMD as a function of centrality are shown in Fig. 3. The $v_n$ values are also obtained by dividing $\chi$ obtained in eqn. 3 above by the fluctuation in the multiplicity in that centrality bin.

The anisotropy parameters for the $\gamma$-like sample are systematically greater than the corresponding parameters for the charged particles and the anisotropy parameters for both of them decrease monotonically with increasing centrality.




Figure 2: The event plane correction factors for charged particles and for $\gamma$-like for the different orders of anisotropy




Figure 3: $v_n$ for different centralities for $\gamma$-like and for charged particles

### 3.3 Decay Effect

The decay of the neutral pions into photons introduces correlations because of energy momentum conservation, causing an apparent anisotropy in the photons which is different from the actual anisotropy present in the pions. The process of decay also dilutes the initial correlation present in the neutral pions. The resultant effect of these two competent effects scales with the experimentally measured quantity $\chi$ and enables a determination of the neutral pion anisotropy using the relation 4 as obtained from reference [3]

$$\frac{v(\gamma)}{v^{in}(\pi^0)} = \frac{a}{(\chi - b)^2} + c \tag{4}$$

The values of the constants in this relation depend weakly on the details of the realistic kinematic distributions of the charged pions. For the present work the values are taken from the reference cited above. For first and second order, the values of (a,b,c) are respectively (0.075, 0.473, 0.801) and (0.031, 0.286, 0.619).

Using the measured values of $v_1$ in the PMD, and using relation 4, the $v_1$ values for neutral pions are obtained, enabling a comparison with the corresponding values for charged particles. The anisotropy values for charged particles, for $\gamma$-like in the PMD and the deduced anisotropy for neutral pions are all shown in Fig. 4. Where this agreement is quite good, there are limitations. Certain effects not incorporated are: the effect of finite granularity on the measured values of anisotropy, the effect of charged particle contamination in the $\gamma$-like sample, the effect of protons on the SPMD and other effects.

### 3.4 Granularity,Efficiency and Other Effects

The PMD has very fine azimuthal granularity, and the finite granularity does not have any effect on the estimated anisotropy values. The azimuthal bin size of the SPMD is 2 degrees. This not-so-fine granularity results in a measured charged particle yield which is less than the actual multiplicity. Since the number of 'hits' and their distribution in SPMD are used, the 'coarse' granularity also has an effect of diluting the actual azimuthal correlation. Events are simulated with initial anisotropy values and multiplicity. The granularity of the SPMD is incorporated before analysing the data for anisotropy. The initial conditions are varied to obtain the experimentally measured values of the anisotropy parameters for charged particles. Using this anisotropy value along with the corresponding multiplicity for neutral pions, followed by neutral pion decay kinematics enable to estimate the anisotropy in photons. These




Figure 4: $v_1$ for different centralities for $\gamma$-like as obtained from the PMD and for charged particles from SPMD. Using $v_1$ from PMD and the scaling relation, $v_1$ for charged particles is estimated.

estimates of anisotropy are compared to those measured in the PMD, for both orders.

The anisotropy measured in the PMD are observed to be larger than what is obtained by the method described above. Further, 60% of the $\gamma$-like sample is assumed to be decay photons from these neutral pions and 40% of the sample is contaminants. We assume that these contaminants carry the same anisotropy as those of the charged particles. We see broad agreement for the first order ($v_1$) but for the second order, $v_2$ observed in PMD is systematically larger than what can be obtained by using the values of $v_2$ for charged particles and using decay kinematics. The results are shown in Fig. 5. These results are preliminary, and at present there are no known explanations for this excess.




Figure 5: $v_n$ for different centralities for $\gamma$-like as obtained from the PMD. Using $v_n$ values of charged particles and decay kinematics, expected values for photon samples are also plotted for two different combinations of efficiency and purity

## 3.5 Event Plane from PMD

Independent of asymmetries in particle emission as studied above, it is possible to determine an event plane from the PMD and investigate any possible asymmetries in charged particle emission, with respect to the event plane. By construction, the asymmetry coefficients (Fourier coefficients) obtained earlier will result in the maximum value. The event plane obtained from the PMD may have a little bias as compared to the event plane for the charged particles due to decay. Event planes estimated for different particle species in different rapidity regions could all be different, beyond fluctuations, the difference arising due to different final state effect on the particles while they stream from the point of freezeout to the detectors. It is then expected that $v_2$ for charged particles with respect to event plane of the PMD would be less than $v_2$ as measured using the event plane from the SPMD. The same result is expected if for the estimate of $v_2$ for $\gamma$-like using the event plane of the SPMD. This is observed and is shown in Fig. 6. This demonstrates that the event plane estimated from the PMD can be used to study emission properties of any particle species, in any other acceptance region.




Figure 6: $v_n$ for different centralities for $\gamma$-like with respect to the event plane obtained in the spmd and for charged particles with respect to event plane obtained in the pmd

### 3.6 Correlation Between Event Planes of Different Orders

A priori, one expects that the event planes of both the orders are either the same, or perpendicular to each other, depending upon the speed of sound in the colliding matter and the colliding energy of the nuclei. The estimates of these event planes are affected with fluctuations.

However, correcting for these fluctuations should result in the same numerical value of $v_2$ determined from event plane of either order. The data from the PMD suggests that this is not the case, where $v_2$ determined from the first order event plane from the PMD is systematically lower, in addition to having larger error bars. The results are shown in Fig. 7a. This indicates that one has to look for possible correlations (or lack of correlations) between the event plane of the two orders. The event planes of the two orders may not be correlated due to effects such as initial state density fluctuations. The absence of this correlation can also arise if the event plane is estimated using particles which are products of decay. Then decay kinematics affects the event planes of different orders differently. To look for correlations between the event plane angles, the quantity $\langle \cos 2(\psi_1 - \psi_2) \rangle$ is deduced, where the average is over the event samples in question. This is plotted as a function of centrality and is shown in Fig. 7b. This behaviour is explicable by the decay effect. Starting with the same direction for the first and the second order event plane, and the different centralities characterised by different mean multiplicities, a simulation which includes the neutral pion decay is able to reproduce the observed correlation between the event planes of the two orders. This shows that the 'primordial' event plane is affected by the decay of the particles, and the quantitative effect is different for the event planes of the two orders. This results into different values of $v_2$ as determined from the event planes of the two orders.

## 4 PMD in the ALICE Experiment

A logical extension of the results obtained from PMD in WA98 experiment was to investigate whether the PMD could measure azimuthal anisotropies in photon emission in the ALICE experiment. The results of these investigations are detailed in reference [11]. Further, can the PMD provide an event plane estimate in the ALICE environment and if this event plane may be used for studying asymmetries in particle emission. This was explored using fast simulation techniques, is detailed in reference [4], and we discuss it here.

The Physics motivation comes from the suggestion that the absorption of




Figure 7: (a) $v_2$ for different centralities for $\gamma$-like with respect to the event plane of first order and event plane of second order. (b) The correlation between the first order and the second order event planes in the PMD

J/$\psi$ by comovers is predicted to be azimuthally asymmetric as compared to an isotropic Glauber absorption in the forward region [9]. The asymmetry has been predicted to be about 5 - 20 % depending upon the centrality, size and the energy of the colliding nuclei. Any observation of this asymmetry in the yield of J/$\psi$ will enable a determination of the relative contribution of the two absorption processes, essential to make an estimate of the actual suppression of the resonance from the possible Quark Gluon Plasma.

The ALICE experiment proposes to measure the muons in the DiMuon Spectrometer D$\mu$S. The PMD is designed to take data at the highest rate possible in ALICE, combining the trigger rates for TPC and D$\mu$S. This makes it a specially attractive detector to provide the event plane necessary for the study of anisotropy in J/$\psi$ absorption. The possibility of measuring any such anisotropy in the J/$\psi$ emission with respect to the event plane measured from the PMD is investigated.

## 4.1 Methodology

The data for the investigation is simulated. The transverse momentum and pseudorapidity distribution of the pions and the kaons are generated by parametrising HIJING events. These are generated in the $\eta$ range 0 to 6 and $p_T$ range 0 to 8 GeV/c. The central events correspond to a maximum charged particle pseudorapidity density of about 6000. A branching ratio of 11.7 mb is assumed for J/$\psi$ decaying into two muons. The yield of muons from other sources is not known a priori, and therefore this number is a free parameter. The muons are added in the acceptance region of the D$\mu$S and their number is proportional to the charged particle multiplicity in the event. The transverse momentum

distribution of these muons was motivated by the distribution of the average number of muons per event greater than a particular $p_T$, as in reference [10].

The azimuthal angles of all the pions and kaons are generated with a certain initial anisotropy. The azimuthal angles of the J/$\psi$s are generated assuming the absorption is maximum along the event plane. This produces a maximum yield of the J/$\psi$ in a direction normal to the event plane determined from the charged particles or from the photons.

## 4.2 Detector Effects

The hits on the PMD arise from both photons and charged hadrons. After filtering hadrons using appropriate algorithms, the remaining hits, called "$\gamma$-like", contain 60% decay photons. The azimuthal distributions of these photons reflect the anisotropy present in parent $\pi^0$. Of the 40% contaminants, half are assumed to be initial charged pions, which have the same anisotropy as the neutral pions, and the remaining half are secondaries likely to be randomly distributed. The generated events contain $\gamma$-like hits having suitable mix of charged particles and decay photons according to the above proportion.

The effect of upstream material, which results in scattering of the incident particles, has been investigated for the coverage of the PMD in Ref. [11]. This effect is incorporated in the fast simulation by allowing all particles to change direction so that the final $\eta - \phi$ differs from the original values by an amount $\delta\eta$ (=0.1,0.2) ,$\delta\phi(= 10^{\circ}, 20^{\circ})$ given by a Gaussian distribution. The smaller values are termed as moderate scattering and the larger values correspond to what we term as large scattering.

Muon track reconstruction efficiency is assumed to be 0.9 in the D$\mu$S. The efficiency has been varied between 0.8 to 1.0 and its effect on the sensitivity of the analysis is also studied. The mass resolution of the J/$\psi$ is considered to be 60 MeV.

## 4.3 Analysis and Results

The invariant mass spectra of opposite sign dimuons is constructed using muons with $p_T > 1$ GeV and the two muons following $p_T^{min} + 0.5\ p_T^{max} > 2$ GeV. To extract the resonance yield, the invariant mass spectra of the like sign dimuons is also obtained with the same kinematic cuts. The distribution is then fitted to a polynomial in the region of the resonance mass. The opposite sign dimuon spectra is then fitted to this polynomial and a Gaussian and the yield of the resonance is determined as shown in Fig. 8




Figure 8: The invariant mass spectra of the opposite sign and the like sign dimuons. The fits to the background and the signal.




Figure 9: S/B ratio and Significance as a function of number of added muons in the D$\mu$S

Since the number of added muons is a free parameter in the simulation, the results for the S/B ratio and the significance of the data sample vary with the number of added muons. The results are shown in Fig. 9.

The event planes of both orders are obtained for every event and the resolution correction factor to correct for fluctuations of the event plane due to finite particle multiplicity. To look for any possible anisotropy in the background spectra, the invariant mass distribution of the like sign muons is plotted in different azimuthal bins with respect to the event plane, and is shown in Fig. 10a. The yield in different azimuthal regions in a fixed mass bin is used for obtaining the anisotropy parameters. This is repeated for different (overlapping) mass bins and the anisotropy parameters determined. There is no measurable anisotropy in the background spectra in the region around the resonance mass, as is shown in Fig. 10b.




Figure 10: (a) Invariant mass spectra of the like sign muons in the different azimuthal bins with respect to the second order event plane from the PMD. (b) Fourier analysis of background yield with respect to the event planes of first and second order in the PMD.

The invariant mass spectra in different azimuthal regions is shown in Fig.11 and the J/$\psi$ yield is then analysed to extract the anisotropy components that describe the azimuthal distribution. This has been repeated for the variety of cases where the particles incident on the PMD remain unscattered by the upstream material, undergo moderate scattering or undergo large scattering.

The observation of anisotropy in J/$\psi$ emission depends on several factors, e.g., background subtraction in the invariant mass spectra, increase in background due to charm and beauty decays, track reconstruction efficiency for muons in the DiMuon Spectrometer and the yield of J/$\psi$ from higher res-




Figure 11: Invariant mass spectra of opposite sign muons in the different azimuthal bins with respect to the second order event plane from the PMD.

onances. A systematic study of variation in these parameters suggests that the *significance* of the data (and hence the number of events required) follows a simple relation with the anisotropy values to be studied and the associated accuracy, and is shown in Fig.12. The fitted curve is *Significance = 0.7/$v_2 \cdot f$* where $v_2$ is the value of anisotropy to be probed and f is the fractional error.




Figure 12: (a) Significance as a function of $v_2 \cdot f$ where $v_2$ is the anisotropy to be probed with a fractional error $f$.(b) Same as (a) without the error due to event plane determination. The fitted curve corresponds to the **minimum** Significance required.

For a given S/B ratio and muon track reconstruction efficiency, this significance can be translated into the number of events required to enable such a determination.

# 5 Summary

The photon multiplicity detector has been successfully implemented in the WA93 and WA98 experiments. The first evidence of azimuthal anisotropy at the SPS energies was reported using measurements from the PMD. Subsequently, PMD has obtained systematic dependence of anisotropy coefficients on centrality and the results have been compared with corresponding results from charged particles. We observe that the elliptic flow observed in the photons is greater than the one expected by assuming same flow for neutral and charged particles. The anisotropy analysis techniques are such that the error on the estimated anisotropy coefficients are likely to decrease with increasing multiplicity and increasing anisotropy. The PMD at STAR and at the ALICE experiment will be able to provide more precise estimates.We have also shown that the event plane from the PMD can be used to study the anisotropic characteristics of particle emission in the ALICE experiment at the LHC.

# 6 Acknowledgements

I wish to thank all the collaborators for help and discussion at various stages of this work and to Dr.Y.P.Viyogi for a critical reading of the manuscript. Part of the work was done at CERN under the CERN-ASIA programme and the CERN Associateship programme. Financial support from the Department of Science and Technology of the Govt.of India is gratefully acknowledged.

# References

[1] M.M.Aggarwal et.al., Nucl.Instr. and Meth.in Phys.Res.**A372** (1996) 143.

[2] M.M.Aggarwal et.al., (WA93 Collaboration), Phys.Lett. **B403** (1997) 390

[3] Rashmi Raniwala, Sudhir Raniwala, Yogendra Pathak Viyogi, Phys.Lett.**B489** (2000) 9

[4] Andreas Morsch, Rashmi Raniwala, Sudhir Raniwala, Yogendra Pathak Viyogi **ALICE-INT-2001-22** (2001)

[5] M.M.Aggarwal et.al., Nucl.Instr. and Meth.in Phys.Res.**A424** (1999) 395.

[6] Bedangadas Mohanty, Invited Talk in ICPAQGP-2001, present proceedings

[7] A.M.Poskanzer, S.Voloshin, Phys.Rev.**C58**(1998)1671

[8] J.-Y.Ollitrault, e-print nucl-ex/**9711003** (1997)

[9] H.Heiselberg, R.Mattielo, Phys.Rev. **C60** (1999) 44902

[10] ALICE Technical Design Report(TDR) of Dimuon Spectrometer, CERN/LHCC **99-22** 1999

[11] ALICE Technical Design Report (TDR) of Photon Multiplicity Detector, CERN/LHCC **99-32** (1999).